

## References

1. Bhutta ZA, Das JK, Bahl R, et al. What will it take to avert preventable newborn deaths and stillbirths and at what cost? *Lancet* 2014; 384: 347–70.
2. Vogel JP, Souza JP, Mori R, et al, and the WHO Multicountry Survey on Maternal and Newborn Health Research Network. Maternal complications and perinatal mortality: findings of the World Health Organization Multicountry Survey on Maternal and Newborn Health. *BJOG* 2014; 121 (suppl 1): 76–88.
3. Macdorman MF, Gregory ECW. Fetal and perinatal mortality, United States, 2013. National vital statistics reports; vol 64 no 8. Hyattsville, MD: National Center for Health Statistics. 2015.
4. Gardosi J, Madurasinghe V, Williams M, Malik A, Francis A. Maternal and fetal risk factors for stillbirth: population based study. *BMJ*. 2013 Jan 24; 346: f108.
5. Chung ST, Sumner AE. Diabetes: T2DM risk prediction in populations of African descent. *Nat Rev Endocrinol*. 2016 Jan 22.
6. Wang S, Alvero R. Racial and ethnic differences in physiology and clinical symptoms of polycystic ovary syndrome. *Semin Reprod Med*. 2013 Sep;31(5):365-9.
7. Jacoby VL, Fujimoto VY, Giudice LC, Kuppermann M, Washington AE. Racial and ethnic disparities in benign gynecologic conditions and associated surgeries. *Am J Obstet Gynecol*. 2010 Jun;202(6):514-21.
8. Borghese B, Sibiude J, Santulli P, Lafay Pillet MC, Marcellin L, Brosens I, Chapron C. Low birth weight is strongly associated with the risk of deep infiltrating endometriosis: results of a 743 case-control study. *PLoS One*. 2015 Feb 13;10(2): e0117387.
9. Doherty DA, Newnham JP, Bower C, Hart R. Implications of polycystic ovary syndrome for pregnancy and for the health of offspring. *Obstet Gynecol*. 2015 Jun;125(6):1397-406.
10. Chu S, Kim S, Lau J, et al. Maternal Obesity and risk of stillbirth: a meta-analysis. *Am J Obstet Gynecol*. 2007; 197: 223-228.
11. Villamor E, Cnattingius S. Interpregnancy weight change and risk of adverse pregnancy outcomes: a population-based study. *Lancet*. 2006; 368:1164–1170.
12. Kristensen J, Vestergaard M, Wisborg K, Kesmodel U, Secher NJ. Prepregnancy weight and the risk of stillbirth and neonatal death. *BJOG* 2005; 112:403–8.
13. Tennant PW, Rankin J, Bell R. Maternal body mass index and the risk of fetal and infant death: a cohort study from the north of England. *Hum Reprod* 2011; 26:1501–11.
14. Huang DY, Usher RH, Kramer MS, Yang H, Morin L, Fretts RC. Determinants of unexplained antepartum fetal deaths. *Obstet Gynecol* 2000; 95:215-21.
15. Rahman FZ, Rahman J, Al-Suleiman SA, Rahman MS. Pregnancy outcome in lupus nephropathy. *Obstet Gynecol Surv* 2004; 59:754-5.
16. Breur J, Kapusta L, Stoutenbeek P, et al. Isolated congenital atrioventricular block diagnosed in utero: natural history and outcome. *J Matern Fetal Neonatal Med*. 2008; 21:469–476.
17. Levesque K, Morel N, Maltret A, Baron G, Masseau A, Orquevaux P, Piette JC, Barriere F, Le Bidois J, Fermont L, Fain O, Theulin A, Sassolas F, Pezard P, Amoura Z, Guettrot-Imbert G, Le Mercier D, Georjin-Lavialle S, Deligny C, Hachulla E, Mouthon L, Ravaud P, Villain E, Bonnet D, Costedoat-Chalumeau N; “Lupus néonatal” group; Group of collaborators. Description of 214 cases of autoimmune congenital heart block: Results of the French neonatal lupus syndrome. *Autoimmun Rev*. 2015 Dec; 14(12):1154-60.
18. Silver RM, Parker CB, Reddy UM, Goldenberg R, Coustan D, Dudley DJ, Saade GR, Stoll B, Koch MA, Conway D, Bukowski R, Hogue CJ, Pinar H, Moore J, Willinger M, Branch DW. Antiphospholipid antibodies in stillbirth. *Obstet Gynecol*. 2013 Sep; 122(3):641-57.
19. Rackham O, Paize F, Weindling AM: Cause of death in infants of women with pregestational diabetes mellitus and the relationship with glycemic control. *Postgrad Med* 2009, 121:26–32.
20. Persson M, Norman M, Hanson U. Obstetric and perinatal outcomes in type 1 diabetic pregnancies. A large, population-based study. *Diabetes Care*. 2009; 32:2005–2009.
21. Ray JG, Burrows RF, Burrows EA, et al. MOSHIP: McMaster outcome study of hypertension in pregnancy. *Early Hum Dev*. 2001; 64:129–143

22. Wahabi HA, Alzeidan RA, Bawazeer GA, Alansari LA, Esmail SA: Preconception care for diabetic women for improving maternal and fetal outcomes: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2010, 10:63.
23. Edwards A, Springett A, Padfield J, Dorling J, Bugg G, Mansell P. Differences in post-mortem findings after stillbirth in women with and without diabetes. *Diabet Med.* 2013 Oct; 30(10):1219-24.
24. Simpson LL. Maternal medical diseases: risk of antepartum fetal death. *Semin Perinatol.* 2002; 26:42–50.
25. Nielsen HS, Witvliet MD, Steffensen R, et al. The presence of HLA-antibodies in recurrent miscarriage patients is associated with a reduced chance of a live birth. *J Reprod Immunol.* 2010 Jul 3.
26. Nielsen HS, Steffensen R, Varming K, Van Halteren AG, Spierings E, Ryder LP, Goulmy E, Christiansen OB. Association of HY-restricting HLA class II alleles with pregnancy outcome in patients with recurrent miscarriage subsequent to a firstborn boy. *Hum Mol Genet.* 2009 May 1; 18 (9):1684-91.